

# Isolation and Screening of Protease Producing Bacteria from Local Environment for Detergent Additive

**Temam Abrar Hamza**

Department of Biotechnology, College of Natural Sciences, Arba Minch University, Arba Minch, Ethiopia

**Email address:**

temam2abrar2@gmail.com

**To cite this article:**

Temam Abrar Hamza. Isolation and Screening of Protease Producing Bacteria from Local Environment for Detergent Additive. *American Journal of Life Sciences*. Vol. 5, No. 5, 2017, pp. 116-124. doi: 10.11648/j.ajls.20170505.11

**Received:** June 15, 2017; **Accepted:** June 23, 2017; **Published:** August 30, 2017

---

**Abstract:** Proteases are among the most important hydrolytic enzymes that found in every organism to undertake important physiological functions. They are multipurpose enzymes used in various industries such as detergent, silver recovery, food, pharmaceutical, leather, and textile industries. This work aimed to produce protease from indigenous microbes for use as detergent additive. Isolation of protease producer was undertaken using skim milk agar medium. Crude enzyme was characterized in terms of wash and stain removal tests. A total of 188 protease positive bacteria were isolated from seven samples collected from Arba Minch town. Out of 36 alkaline protease producing bacteria, one isolate designated as *Bacillus sp. Cab44* was selected. The optimum activity was reached at pH 9 and 50°C. The enzyme was stable in the pH range of 7 to 10. It retained 75%, 86% and 72% activity after one hr pre-incubation at 50°C, in 15% H<sub>2</sub>O<sub>2</sub> and 0.3% commercial detergent respectively. The enzyme activity was increased by Mg<sup>2+</sup>, Cu<sup>2+</sup>, and Mn<sup>2+</sup>, was not affected by Ca<sup>2+</sup> but decreased by Zn<sup>2+</sup>, Hg<sup>2+</sup> and Fe<sup>2+</sup>. It removed stains of blood and egg yolk on cotton cloth at pH 9, 40°C, 5.07 U/ml in 30-40 min. These properties suggest that protease from *Bacillus sp. Cab44* could find potential application in detergent industries as good candidate of additive in detergent formulation which have economic implication.

**Keywords:** *Bacillus sp. Cab44*, Detergent Additive, Enzyme, Protease

---

## 1. Introduction

Proteases are among the most important hydrolytic group of enzymes that found in every organism (prokaryotes, fungi, plants and animals) to undertake important physiological functions [1]. Modern science (microbial biotechnology) explores and exploits the beneficial microbial wealth for various human requirements. Humans over the ages have been highly successful in applying processes carried out by microorganisms to solve problem in various industry and environmental quality. Although, animal and plant proteases are of important industrial applications, a large proportion of commercially useful proteolytic enzymes currently available in the market are from microorganisms [2].

Microbial proteases are one of the most important groups of enzymes, used in various industrial processes as food, pharmaceutical and detergent industries, as well as in the preparation of leather, textile and wool, among others [3, 4, 5, 6]. It has also promising application in medical usage and management of industrial and household waste. The use of

microbial system (enzymes) is the best alternative for generation of pollution free industries.

The idea of using detergent enzymes dates back to 1914 when two German scientists, Rohm and Haas, used pancreatic proteases and sodium carbonate in washing detergents. However, it was only in 1963 alkaline protease was effectively incorporated in detergent powder. Due to this its economic importance became well known [7]. However, currently proteases constitute the largest product segment in the global industrial enzymes market. The detergent enzyme market has grown nearly 10-fold during the past 20 years [8].

Removal of proteinaceous stains such as blood, milk, egg, grass and chocolate is very difficult using conventional detergent method. However, removal of such stain is achieved by using alkaline proteases [9, 10, 11]. In addition the use of protease supplementation to detergent formulation significantly improves the cleansing of proteinaceous stain and to deliver unique benefits that cannot otherwise be obtained with conventional detergent technologies [12, 13].

Over the past 30 years, the proteases in detergents have changed from being minor additives to being the key ingredients.

Most commercially available detergents have an alkaline pH and contain chelating agents to overcome water hardness. As result enzyme used for detergent application need to be active and stable at alkaline pH, ionic strength of detergent solution, different washing temperature range (25-60°C). There are also many more parameters involved in the selection of a good detergent protease, such as compatibility with detergent components, e.g. surfactants, perfumes and bleaches; stain degradation and shelf life [3, 7]. Good detergent enzyme should also be stable in the presence of oxidizing agents and bleaches.

Despite the vast microbial diversity of Ethiopia, alkaline protease producing bacteria have not yet been explored. Therefore a research project has been initiated to investigate alkaline protease from bacteria isolated from local habitat and examine their potential to be used as detergent additives.

## 2. Materials and Methods

### 2.1. Materials

The major instruments used in this study were Spectrophotometer, incubator, autoclave, micropipette, rotary shaker and water bath shaker. The major chemicals used were Folin Ciocalteu's phenol reagent, different buffers, hydrogen peroxide, HCl NaOH, Casein and trichloroacetic acid.

### 2.2. Methods

#### 2.2.1. Sample Collection and Isolation of Alkalophilic and Proteolytic Bacteria

Samples were collected from different environments around Arba Minch like abattoir, waste disposing area, fish processing area, compost processing site, Abaya and Chamo Lakes. Each sample was kept in clean sterile sample bottles sealed and transferred to the laboratory and stored at 7°C.

The soil and water samples were suspended in water by vigorous vortexing and serial dilutions were made up to  $10^{-6}$  in sterile distilled water. 0.1 ml of appropriate dilution were added to petri plate on skim milk agar plate containing peptone (0.1%), NaCl (0.5%), Agar (2%) and skimmed milk (10%) at pH 9 and incubated at 40°C for three days. Alkaliphilic proteolytic bacteria were screened. A clear zone formed colonies at pH 9 were picked and purified by streaking on skim milk agar. The cultures were subsequently sub-cultured and used regularly. Agar slants were prepared and preserved at 7°C for further experiments and in 25% glycerol stocks at -25°C for long term storage.

#### 2.2.2. Screening of Potential Isolates

Five alkalophilic strains were screened on the basis of clear zone size for further selection of single isolate on the quality of enzyme produced under submerged condition. The supernatant from each isolate was used as crude enzyme for

observing individual wash performance. Isolate which had high potential on egg stain removal efficiency were selected for further characterization. Finally colonial morphology, cellular morphology and biochemical characteristics were undertaken. Identification of the selected potential isolate was done according to Bergey's manual of determinative bacteriology. Colony morphology characterized by direct observation of the overnight grown microorganisms on skim milk agar plate.

#### 2.2.3. Production of Protease, Protease Assay and Enzyme Unit (U/ml)

Protease production media containing (g/l): Glucose 10, Peptone 7.5,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  5,  $\text{K}_2\text{HPO}_4$  2, CaCl 1 [14] was prepared for enzyme production. The pH of the medium was adjusted to 9. The bacterial isolate was inoculated with 1% of inoculum into 250 ml Erlenmeyer flasks containing 100 ml production medium followed by incubation at 30°C for 48 hours in rotary shaker at 150 rpm. Culture filtrates were separated by centrifugation at 5000 rpm for 15 min and the supernatants were used as crude enzyme source for quantitative enzyme assay.

Protease activity in the culture supernatant was determined using casein as a substrate with some modification of the method described by Gaur *et al.* (2010) [15]. The enzyme activity was expressed in units (U) and calculated by using tyrosine standard calibration curve. One unit of enzyme was defined as the amount of enzyme that releases 1  $\mu\text{mol}$  of tyrosine per ml of crude extract per minute under standard conditions at 40°C. The tyrosine standard curve was used to generate the enzyme unit determination formula in U/ml.

#### 2.2.4. Characterization of Crude Enzyme

The protease activity of crude enzyme at different pH value was tested with 0.005M sodium bicarbonate buffer (pH 7.0 to 10.0) and 0.02M potassium chloride/NaOH buffer (pH 11.0 to 12.0). pH stability was determined by preincubating the crude supernatant in buffers of pH (7-12) for 1 hr. The enzyme activity was measured at 40°C.

Temperature effect on protease activity was determined at different temperatures (30-70°C) [16]. To determine thermal stability, the samples were pre-incubated at different temperatures (45 and 50°C) following determination of residual activity every 20 minutes. Relative protease activity (expressed in%) was defined as the percent protease activity compared with the maximum value.

To investigate the effect of oxidizing agents on enzyme stability, hydrogen peroxide was used from 0-30% in the reaction mixture. Pre-incubation of the reaction mixtures was carried out by mixing equal amount of enzyme and  $\text{H}_2\text{O}_2$  [17].

To determine the effect of metal ions on enzyme activity, the enzyme assay was carried out in the presence of 0.01M  $\text{CaCl}_2$ ,  $\text{MgSO}_4$ ,  $\text{FeSO}_4$ ,  $\text{ZnCl}_2$ ,  $\text{MnSO}_4$ ,  $\text{HgCl}_2$  and  $\text{CuSO}_4$ . The activity of the enzyme without any additives was taken as 100%.

### 2.2.5. Stain Removal and Detergent Compatibility of Crude Enzyme

A commercial powder detergent, Aerial was used for detergent compatibility test of enzyme produced by the selected isolate. The detergent was heated at 100°C for 1hr to denature all enzymes found in the detergent [18]. Crude enzyme was preincubated with 0.3% (w/v) heated detergent for 1 hr and residual activity was measured to check the compatibility of the enzyme with all the surfactants found in the detergent. Enzyme without detergent was used as a control.

Wash test with protease preparation: Wash performance of crude enzyme was determined according to the method of Pathak and Deshmukh (2012) [18] with little modification. White test fabric (cotton) cloth piece (5×5 cm) stained with animal blood or egg yolk was allowed to dry in open air for 5 hr. The stained cloth was put in petri dish and treated with 10 ml crude enzyme at 40°C for 30-40 min. Phosphate buffer was used as control under the same condition. Protein stain removal was checked qualitatively by visualization.

### 2.3. Experimental Design and Statistical Analysis

Average values of duplicate experiments were taken. Microsoft office Excel worksheet 2010 was used for data analysis and presentation.

## 3. Results

### 3.1. Isolation of Alkalophilic and Proteolytic Bacteria

One hundred eighty eight protease positive isolates were obtained from seven sample areas 61, 20, 11, 38, 22, 13 and 23 from Abaya Lake, Chamo Lake, AMU soil, compost processing site, abattoir, waste disposing area and fish processing area respectively. Hence the strains were identified as a protease producer (Figure 1). Out of 188, 36 alkaline protease positive bacterial isolates were screened and five isolates with relatively higher clear zones were further examined.



**Figure 1.** Zone of hydrolysis by strain Cab44 in milk agar plate after 24 h incubation at 40°C.

Among the five isolates, one potential isolate with high egg yolk stain removal efficiency from cloths were selected for further characterization. The results are summarized in Table 1.

**Table 1.** Results of stain removal efficiency of five selected isolates.

Sample Area	Strain code	Egg yolk stain removal efficiency	Remark
-	Control		Before Treatment
AMU Soil	Cam5		After Treatment
Fish Processing Area	Cfp10		
Abaya Lake	Cal16		
Chamo Lake	Ccl33		
Abattoir	Cab44		

### 3.2. Morphological and Biochemical Characteristics of Isolate Cab44

Table 2 shows results of morphological and biochemical tests of the selected isolate. The isolate was characterized as Gram positive and catalase positive. The colonies were

characterized as opaque, irregular and spready configuration and irregular margin. Motile, long irregular and rod shaped cells were observed under light microscope with 1000X magnification. The strain was able to hydrolyse casein. Based on this the isolate was designated as *Bacillus sp. Cab44*.




Figure 2. Gram reaction result (+ve) (a) and Endospore of *Bacillus sp. Cab44* (b) at 40°C and 24 h on milk agar.




Figure 3. Effect of temperature on enzyme activity after 30 min incubation at different temperature.

Table 2. Results of the morphological and biochemical tests of isolate Cab44.

Morphological characteristics	Results
Colour	Creamy white
Size	Large
Margin	Irregular
Elevation	Flat
Configuration	Irregular and spread
Morphology	Rod
Optical density	Opaque
Gram reaction	Positive
Endospore	Elliptical
Motility test	positive
Biochemical and physiological characteristics	Results
Nitrate reduction	Negative
Indole tests	Positive
MR-VP test	Negative
Citrate utilization	Negative
Casein hydrolysis	Positive

### 3.3. Characterization of Crude Enzyme


#### 3.3.1. Effect of Temperature on Enzyme Activity and Stability

The protease produced from *Bacillus sp. Cab44* was found to be active at a range of temperatures, between 30 and 60°C. Figure 3 shows that maximum enzyme activity was recorded at 50°C and the enzyme retained 16% of its maximum activity at 70°C. Around 80% of its maximum activity was expressed at 40 and 55°C.


The enzyme was stable at 45 and 50°C retaining more than 90% and 75% of its maximum activity for 60 min of pre-incubation respectively.

#### 3.3.2. Effect of pH on Enzyme Activity and Stability


The enzyme was active in the range of pH 7-11 as shown in Figure 5. The optimum activity pH was 9.0. The enzyme showed 89% and 82.1% of its maximum activity at pH 8.0 and 10.0 respectively.



**Figure 4.** Effect of temperature on enzyme stability after 60 min incubation at 45 and 50°C for 20 min interval regularly.



**Figure 5.** Effect of pH on enzyme activity (at 40°C for 30 min incubation) and stability (after 60 min pre-incubation in different pH buffers at 30°C) then assay was performed at 40°C and pH 9. Activity of the enzyme pre-incubated at pH 9 and assayed at same pH was considered as 100%.



**Figure 6.** Effect of hydrogen peroxide on enzyme stability after preincubation for 60 min at 30°C following assay at 40°C and pH of 9.

The enzyme was stable in a broad pH range, maintaining over 83 and 76% of its maximum activity at pH 7 and 10 respectively (Figure 5). However, only 19% of the maximum enzyme activity was retained at pH 12.

### 3.3.3. Effect of Oxidizing Agent (H<sub>2</sub>O<sub>2</sub>) and Detergent on Enzyme Stability

The enzyme retained 85.6% of its activity when pre-incubated in 15% H<sub>2</sub>O<sub>2</sub> for an hour. As the percentage of H<sub>2</sub>O<sub>2</sub> increased the residual activity was decreased with 28%

at 30% H<sub>2</sub>O<sub>2</sub> (Figure 6). The enzyme showed 72% of its activity after 1 h pre-incubation in 0.3% commercial detergent (Ariel).

### 3.3.4. Effect of Metal Ions on Enzyme Activity

The different cations tested at 0.01M concentration affected the enzyme activity differently. The activity was enhanced by Mg<sup>2+</sup> (129%), Cu<sup>2+</sup> (113%) and Mn<sup>2+</sup> (110%). It was inhibited by Zn<sup>2+</sup> (65.6%), Hg<sup>2+</sup> (43.5%) and Fe<sup>2+</sup> (2.2%). However, Ca<sup>2+</sup> did not affect the activity (Figure 7).






Figure 7. Effect of metal ions on enzyme activity for 30 min incubation at 40°C and pH 9.

### 3.4. Stain Removal Efficiency of Crude Enzyme

The crude enzyme produced from *Bacillus sp. Cab44* efficiently removed egg yolk stain in 40 min. It took around 30 min to remove blood stain from cotton cloth (Figure 8 and 9).




Figure 8. Egg yolk stain removal efficiency of enzyme produced by *Bacillus sp. Cab44*: (T) normal cotton Cloth; (A) Stained cotton fabric before reaction; (B) control (0% of enzyme/ only phosphate buffer); (C) with 5.07U/ml of crude enzyme for 40 min at 40°C and pH 9.




Figure 9. Blood stain removal efficiency of enzyme produced by *Bacillus Sp TF44*: (T) normal cotton Cloth; (A) Blood stained cotton fabric before reaction; (B) control (0% of enzyme, only phosphate buffer); (C) with 5.07U/ml of crude enzyme for 30 min at 40°C and pH 9.

## 4. Discussion

One of the main concerns of this study was to isolate and identify alkalophilic isolate having a vital ability to secrete

extra-cellular proteolytic enzyme to be used for detergent additive. Accordingly, 188 bacterial strains were isolated from seven different areas. Formation of clear zone around the bacterial colony indicated the protease positive strains hydrolysed the skim milk present in the media (Figure 1).

Five isolates were screened on the basis of clear zone size. Out of these, *Bacillus sp. Cab44* exhibited high capability of removing egg yolk stain compared to others (Table 1). The use of alkaline skim milk agar for the isolation of alkaline protease producing bacteria has earlier been reported by some workers [19, 20]. Results show that, Abaya Lake is rich in proteolytic bacteria. This indicates that Abaya Lake has potential for screening of bacterial strain which produce industrially important enzymes.

The crude enzyme produced from *Bacillus sp. Cab44* was further characterized to evaluate its potential applications for formulation of detergent. It was optimally active between pH 7-10 and stable at alkaline pH. Proteases active in the pH range of 8-12 and stable at alkaline pH are known as potential candidates for detergent application. This indicates that protease produced from *Bacillus sp. Cab44* has a good potential for such applications [21, 22, 23, 24].

Another interesting property of the protease produced from *Bacillus sp. Cab44* was its ability to maintain more than 75% of its maximum activity after pre-incubation at its optimum temperature (50°C) for 60 min. The studies on temperature profile of the enzyme showed an optimum temperature range between 40-60°C with maximal activity at 50°C (Figure 3). The physiochemical characterization of the protease indicates its suitability as a potential additive in detergents [18, 19, 23, 24]. These observations were also equivalent to that had been found by Jabeen and Qazi (2011) [25]. Maximum activity of protease produced from *Bacillus sp. Cab44* at high pH (9) and temperature of 50°C indicates its high potential of industrial employment in hard wash condition.

The stability and compatibility of produced enzyme with oxidizing agent is an important factor to be used as detergent additive. Detergents contain oxidizing agents, which may hinder the activity of certain enzymes and hence enzyme stable with oxidizing agents are required for detergent industry. The protease produced from the isolate lost little or no enzyme activity on treatment with 15% H<sub>2</sub>O<sub>2</sub> for one hour. This shows that the protease produced from the isolate is compatible with and stable in oxidizing agents [26].

Identification of appropriate metal ions for activity and thermo stability to the enzymes are very important for their applications at commercial levels [27]. Among the tested metal ions Mg<sup>+2</sup> is potent inducer of the protease activity as shown in Figure 7. On the other hand Hg<sup>+2</sup> and Fe<sup>+2</sup> are the potent inhibitors of enzymatic activity produced from *Bacillus sp. Cab44*. Maximum loss of enzymatic activity was observed in Fe<sup>+2</sup>. The presence of Ca<sup>2+</sup> did not affect the enzyme activity [28]. However, in another report, metal ions like Mn<sup>2+</sup> and Ca<sup>2+</sup> were found to be potent enhancers [29].

In the present investigation (shown in Figure 8 and 9) the protease produced from *Bacillus sp. Cab44* had high capability of removing blood and egg yolk stains. Interestingly, the stain removal process was accomplished without any commercial detergent additives. This makes the protease advantageous for use as detergent additive against a wide variety of stains like blood, egg yolk and other body secretions from cloths. Also it was compatible with all

surfactants found in commercial detergent (Ariel), which indicates its potential candidate in detergent formulation. This result is well established by many authors [25, 28, 30]. Such practice may find application in removing blood stains from materials of medical importance. The result is far better comparable to the earlier reports of protease from *Pseudomonas aeruginosa* [9] and *Bacillus licheniformis* U1 [18].

The enzyme used in this study has the ability to completely remove blood and egg yolk stains within 40 min at 40°C at 5.07 U/ml, unlike the protease that was used to remove egg yolk stain from test fabric which needed 2h for complete egg yolk stain removal at 40°C and much higher enzyme concentration (100U/ml) [31].

The exploration of protease producing bacteria is still underway to achieve high quality industrial grade enzyme to produce goods [32]. The use of microorganisms to produce enzymes has a number of technical and economic advantages and in recent years it has become the predominant mode of enzyme production [33, 34].

## 5. Conclusions

As the method in this investigation, the accomplishment of stain removal process with a minimum time requirement indicates that this enzyme is highly cost effective candidate for enzyme incorporation in detergent formulations. In addition removal of stain from cloths undergoes the process without any additive of detergent or chemicals. These indicate the enzymes produced in this investigation have potential application in stain removal of medical materials.

## Acknowledgements

The author wish to thank Ato Zerihun Senbeto for his financial support for publication of this research paper. Also the author would like to thank to Zemzem Ayele, Haniza Redwan and Omer Abrar for their supports.

## References

- [1] Rao, M. B., Tanksale, A. M., Ghatge, M. S. and Deshpande, V. V. (1998). Molecular and biotechnological aspects of microbial proteases. *Microbiology and Molecular Biology Research*, 62:597-635.
- [2] Saperas, N. and Subir, F. E. (2011). Proteolytic enzymes in detergents: evidence of their presence through activity measurements based on electrophoresis. *Journal of Chemistry*, 88:1702-1706.
- [3] Gupta, R., Beg, Q. K. and Lorenz, P. (2002). Bacterial alkaline proteases: molecular approaches and industrial applications. *Applied Microbial Biotechnology*, 59:15-32.
- [4] Saha, M. L., Begum, K. J. M. H., Khan, M. R. and Gomes, D. J. (2011). Bacteria associated with the tannery effluent and their alkaline protease activities. *Plant Tissue Culture Biotechnology*, 21:53-61.

- [5] Younes, G., Sara, R., Alireza, E., Aboozar, K., Maryam, S. and Najme, T. (2011). Screening and isolation of extracellular protease producing bacteria from the Maharloo Salt Lake. *Iranian Journal of Pharmaceutical Sciences*, 7:175-180.
- [6] Vadlamani, S. and Parcha, S. R. (2011). Studies on industrially important alkaline protease production from locally isolated superior microbial strain from soil microorganisms. *International Journal of Biotechnology Applications*, 3:102-105.
- [7] Kumar, D., Savitri, T., Verma, R. and Bhalla, T. C. (2008). Microbial proteases and application as laundry detergent additive. *Research Journal of Microbiology*, 3:661-672.
- [8] Rajkumar, R., Jayappriyan, R. K. and Rengasamy, R. (2011). Purification and characterization of a protease produced by *Bacillus megaterium* RRM2: application in detergent and dehairing industries. *Journal of Basic Microbiology*, 51:614-624.
- [9] Najafi, M. F. (2005). Potential application of protease isolated from *Pseudomonas aeruginosa* PD100. *Electronic Journal of Biotechnology*, 8:197-203.
- [10] Raj, A., Khes, N., Pujari, N., Bhattacharya, S., Das, A. and Rajan, S. S. (2012). Enhancement of protease production by *Pseudomonas aeruginosa* isolated from dairy effluent sludge and determination of its fibrinolytic potential. *Asian Pacific Journal of Tropical Biomedicine*, 3:1845-1851.
- [11] Chaudhari, S. G., Chaudhari, S. D., Khobragade, R. M. and Deshmukh, A. M. (2013). Studies on production of alkaline proteases from *Bacillus* species and its application in detergent industry. *Dav International Journal of Science*, 2:44-48.
- [12] Bayoumi, R. A., Atta, M. H., Swelam, S. M. and El-Hemiany, A. (2010). Microbial production of thermo alkaliphilic enzymes for application in biodetergent technology. *European Journal of Applied Sciences*, 2:85-93.
- [13] Prabhavathy, G., Rajasekara, M. P. and Senthilkumar, B. (2013). Identification of industrially important alkaline protease producing *Bacillus subtilis* by 16s rRNA sequence analysis and its applications. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 4:332-338.
- [14] Das, G. and Prasad, M. P. (2010). Isolation, purification & mass production of protease enzyme from *Bacillus subtilis*. *International Research Journals of Microbiology*, 1:26-31.
- [15] Gaur, S., Agrahari, S. and Wadhwa, N. (2010). Purification of protease from *Pseudomonas thermaerum* GW1 isolated from poultry waste site. *The Open Microbiology Journal*, 4:67-74.
- [16] Padmapriya, B., Rajeswari, T., Nandran, R. and Raj, F. (2012). Production and purification of alkaline serine protease from marine *Bacillus* species and its application in detergent industry. *European Journal of Applied Sciences*, 4:21-26.
- [17] Pathak, A. P. and Deshmukh, K. B. (2012). Alkaline protease production, extraction and characterization from alkaliphilic *Bacillus licheniformis* KBDL4: A Lonar Soda Lake isolate. *Indian Journal of Experimental Biology*, 50:569-576.
- [18] Pravin, D., Sunil, B., Anjana, G. and Bhatt, S. (2014). Isolation, characterization and investigating the industrial applications of thermostable and solvent tolerant serine protease from hot spring isolated thermophilic *Bacillus licheniformis* U1. *International Journal of Applied Science and Biotechnology*, 2:75-82.
- [19] Boominadhan, U., Rajakumar, R., Karppaga, P. V. S. and Joe, M. M. (2009). Optimization of protease enzyme production using *Bacillus* sp. Isolated from different wastes. *Botany Research International*, 2:83-87.
- [20] Haile, G. and Gessesse, A. (2012). Properties of alkaline protease C45 produced by alkaliphilic *Bacillus* sp. isolated from Chitu, Ethiopian Soda Lake. *Journal of Biotechnology and Biomatter*, 2: 1-4.
- [21] Rao, K. and Narasu, L. M. (2007). Alkaline protease from *Bacillus firmus* 7728. *African Journal of Biotechnology*, 6:2493-2496.
- [22] Sepahy, A. A. and Jabalameli, J. (2011). Effect of culture conditions on the production of an extracellular protease by *Bacillus* sp. isolated from soil sample of Lavizan Jungle Park. *SAGE-Hindawi Access to Research Enzyme Research*, 1:1-7.
- [23] Verma, T. and Baiswar, V. (2013). Isolation and characterization of extracellular thermoalkaline protease producing *Bacillus cereus* isolated from tannery effluent. *The International Journal of Engineering and Science*, 2:23-29.
- [24] Vijayaraghavan, P., Lazarus, S. and Gnana, S. P. V. (2013). De-hairing protease production by an isolated *Bacillus cereus* strain AT under solid-state fermentation using cow dung: biosynthesis and properties. *Saudi Journal of Biological Sciences*, 1:1-8.
- [25] Jabeen, F. and Qazi, J. I. (2011). Production and optimization of detergent compatible thermostable alkaline protease from *Bacillus cereus* FJ10. *Journal of Scientific and Industrial Research*, 70:1042-1048.
- [26] Chu, W. H. (2007). Optimization of extracellular alkaline protease production from species of *Bacillus*. *Journal of Indian Microbiology and Biotechnology*, 34:241-245.
- [27] Nadeem, M., Qazi, J. I., Baig, S. and Syed, Q. (2007). Studies on commercially important alkaline protease from *Bacillus licheniformis* N-2 isolated from decaying organic soil. *Turkish Journal of Biochemistry*, 32:171-177.
- [28] Rajkumar, R., Jayappriyan, R. K. and Rengasamy, R. (2011). Purification and characterization of a protease produced by *Bacillus megaterium* RRM2: application in detergent and dehairing industries. *Journal of Basic Microbiology*, 51:614-624.
- [29] Siddalingeshwara, K. G., Uday, J., Huchesh, C. H., Puttaraju, H. P., Karthic, J., Sudipta, K. M., Pramod, T. and Vishwanatha, T. (2010). Screening and characterization of protease from *Bacillus* sp. *International Journal of Applied Biology and Pharmaceutical Technology*, 1:575-581.
- [30] Hema, T. A. and Shiny, M. (2012). Production of protease enzyme from *Bacillus clausii* Sm3. *Journal of Pharmacy and Biological Science*, 1:1-37.
- [31] Kumar, D. and Bhalla, T. C. (2004). *Bacillus* sp. APR-4 protease as a laundry additive. *Indian Journal of Biotechnology*, 3:563-567.
- [32] Priya, V. V. G., Preethi, S., Karthikeyan, S. and Babu, R. N. G. (2016). Isolation and identification of protease producing bacteria from soil. *International Journal of Research in Engineering and Technology*, 3 (8):1362-1365.



- [33] Singh, P., Rani, A. and Chaudhary, N. (2015). Isolation and characterization of protease producing *Bacillus* sp from soil. *International Journal of Pharma Sciences and Research*, 6 (4):633-639.
- [34] Harish R. and Chauhan, B. J. (2017). Isolation, characterization of protease producing microbes from soil of agriculture land and purification of protease. *International Journal of Pharma Research and Health Sciences*, 5 (1):1581-1585.